

It is not believed that extensions of time or fees for net addition of claims are required beyond those that may otherwise be provided for in documents accompanying this paper. However, if additional extensions of time are necessary to prevent abandonment of this application, then such extensions of time are hereby petitioned under 37 C.F.R. § 1.136(a), and any fees required therefor (including fees for net addition of claims) are hereby authorized to be charged to our Deposit Account No. 19-0036.

Amendment

In the Specification:

Please substitute the following paragraph for the pending paragraph bridging pages 7 and 8 of the specification:

Figs. 1A-1C depict the nucleotide and translated amino acid sequence (Seq ID Nos. 1 and 2) of the AD7c-NTP cDNA. The shaded region corresponds to the nucleic acid sequences detected in 6 AD brains by RT-PCR analysis of mRNA. The cDNA exhibits significant homology with Alu gene, and to an unknown gene in the Huntington region, Chromosome 4q16.3 (underlined). The open reading frame begins with the first methionine codon. The translated amino acid sequence encodes a 41.3 kD protein with a hydrophobic leader sequence (*italics*) followed by a myristoylation motif (***bold, italics***) and potential AI cleavage site. That same region (*italics, underlined*) exhibits significant

A¹

09964667-092801

homology with the insulin/IGF-1 chimeric receptor. There are 17 potential glycogen synthase kinase-3, protein kinase C, or cAMP or Ca-dependent kinase II phosphorylation motifs and one transforming growth factor (tgf) motif (double underlined). The embolded amino acid sequences exhibit significant homology with the A4 alternatively spliced mutant form of NF2, β subunit of integrin, and human decay accelerating factor 2 precursor. The boxed amino acid sequences exhibit significant homology with human integral membrane protein and myelin oligoglycoprotein-16.

Please substitute the following paragraph for the pending first full paragraph at page 8 of the specification:

Figs. 2A-2F depict AD7c-NTP expression *in vitro* and *in vivo*. (2A):

A²

Recombinant protein detected by *in vitro* translation using sense strand cRNA transcripts. (2B): Western blot analysis of purified recombinant protein demonstrating specific immunoreactivity with the Tag and N3I4 AD7c-NTP monoclonal antibodies, but not with non-relevant FB50 monoclonal antibody. (2C): Western blot analysis of BOSC cells stably transfected with pcDNA3-AD7c-NTP or pcDNA3 (empty vector). The blots were probed with the N3I4 AD7c-NTP antibody. (2D): Significantly increased levels of the 41-45 kD AD7c-NTP protein in AD frontal lobe relative to age-matched control frontal lobe tissue. Similar results were obtained for temporal lobe tissue. (2E): Higher levels of the 41-45 kD and 19-21 kD AD7c-NTP proteins in late, end-stage (L) AD compared with early, less symptomatic (E) AD. All tissue samples were taken from the

Appl. No. To Be Assigned
(Div. of Appl. No. 09/380,203)

A²
frontal lobe. Note the clusters of 3 or 4 bands between ~41 and ~45 kD, probably corresponding to different degrees of phosphorylation. (2F): Western blot analysis of postmortem ventricular fluid demonstrating higher levels of the ~41 kD AD7c-NTP molecules in AD compared with aged control samples using the N3I4 antibody. The ~28-30 kD band may represent a degradation product. Also note detection of the ~19-21 kD N3I4-immunoreactive molecules in AD.

✓ Please cancel the existing Sequence Listing for the above-identified application and replace it with the substitute Sequence Listing appended hereto.

09364637-092804
108260-29949650